

Sequences of Inequalities among Differences of Gini Means and Divergence Measures

Inder Jeet Taneja

Departamento de Matemática
Universidade Federal de Santa Catarina
88.040-900 Florianópolis, SC, Brazil.

e-mail: taneja@mtm.ufsc.br
<http://www.mtm.ufsc.br/~taneja>

Abstract

In 1938, Gini [3] studied a mean having two parameters. Later, many authors studied properties of this mean. In particular, it contains the famous means as harmonic, geometric, arithmetic, etc. Here we considered a sequence of inequalities arising due to particular values of each parameter of Gini's mean. This sequence generates many nonnegative differences. Not all of them are convex. We have studied here convexity of these differences and again established new sequences of inequalities of these differences. Considering in terms of probability distributions these differences, we have made connections with some of well known divergence measures.

Key words: *Arithmetic mean; Geometric Mean; Harmonic Mean; Gini Mean; Power Mean; Differences of Means; Divergence measures*

AMS Classification: 94A17; 26A48; 26D07.

1 Gini Mean of order r and s

The Gini [3] mean of order r and s is given by

$$E_{r,s}(a, b) = \begin{cases} \left(\frac{a^r + b^r}{a^s + b^s} \right)^{\frac{1}{r-s}} & r \neq s \\ \exp \left(\frac{a^r \ln a + b^r \ln b}{a^r + b^r} \right) & r = s \neq 0 \\ \sqrt{ab} & r = s = 0 \end{cases} \quad (1)$$

In particular when $s = 0$ in (1), we have

$$E_{r,0}(a, b) := B_r(a, b) = \begin{cases} \left(\frac{a^r + b^r}{2} \right)^{\frac{1}{r}}, & r \neq 0 \\ \sqrt{ab}, & r = 0 \end{cases} \quad (2)$$

Again, when $s = r - 1$ in (1), we have

$$E_{r,r-1}(a, b) := K_s(a, b) = \frac{a^r + b^r}{a^{r-1} + b^{r-1}}, \quad r \in \mathbb{R} \quad (3)$$

The expression (2) is famous as *mean of order r or power mean*. The expression (3) is known as *Lehmer mean*[4]. Both these means are monotonically increasing in r . Moreover, these two have the following inequality [2] among each other:

$$B_r(a, b) \begin{cases} < L_r(a, b), & r > 1 \\ > L_r(a, b), & r < 1 \end{cases} \quad (4)$$

Since $E_{r,s} = E_{s,r}$, the Gini-mean $E_{r,s}(a, b)$ given by (1) is an increasing function in r or s . Using the monotonicity property [1], [7], [5] we have the following inequalities:

$$\begin{aligned} E_{-3,-2} &\leq E_{-2,-1} \leq E_{-3/2,-1/2} \leq E_{-1,0} \leq E_{-1/2,0} \\ &\leq E_{-1/2,1/2} \leq E_{0,1/2} \leq E_{0,1} \leq E_{0,2} \leq E_{1,2} \end{aligned} \quad (5)$$

or

$$\begin{aligned} E_{-3,-2} &\leq E_{-2,-1} \leq E_{-3/2,-1/2} \leq E_{-1,0} \leq E_{-1/2,0} \\ &\leq E_{-1/2,1/2} \leq E_{0,1/2} \leq E_{0,1} \leq E_{1/2,1} \leq E_{1,2} \end{aligned} \quad (6)$$

Equivalently, let us write as

$$\begin{aligned} K_{-2} &\leq K_{-1} \leq K_{-1/2} \leq (K_0 = B_{-1} = H) \leq B_{-1/2} \\ &\leq (K_{1/2} = B_0 = G) \leq B_{1/2} \leq (K_1 = B_1 = A) \leq (B_2 = S) \leq K_2, \end{aligned} \quad (7)$$

or

$$\begin{aligned} K_{-2} &\leq K_{-1} \leq K_{-1/2} \leq (K_0 = B_{-1} = H) \leq B_{-1/2} \\ &\leq (K_{1/2} = B_0 = G) \leq B_{1/2} \leq (K_1 = B_1 = A) \leq E_{1/2,1} \leq K_2. \end{aligned} \quad (8)$$

where H , G , A and S are respectively, the *harmonic*, *geometric*, *arithmetic* and the *square-root* means. We observe from the expression (7) that the means considered either are the particular cases of (2) or of (3), while in the second case we have we have a mean $E_{1/2,1}$, which is neither a particular case of (2) nor of (3). We can easily find values proving that there is no relation between S and $E_{1/2,1}$. Summarizing the expressions (7) and (8), we can write them in joint form as

$$P_1 \leq P_2 \leq P_3 \leq H \leq P_4 \leq G \leq N_1 \leq A \leq (P_5 \text{ or } S) \leq P_6, \quad (9)$$

where $P_1 = K_{-2}$, $P_2 = K_{-1}$, $P_3 = K_{-1/2}$, $P_4 = B_{-1/2}$, $N_1 = B_{1/2}$, $P_5 = E_{1/2,1}$ and $P_6 = K_2$.

In [9, 10], the author studied the following inequalities:

$$H \leq G \leq N_1 \leq N_3 \leq N_2 \leq A \leq S, \quad (10)$$

where

$$N_2(a, b) = \left(\frac{\sqrt{a} + \sqrt{b}}{2} \right) \left(\sqrt{\frac{a+b}{2}} \right)$$

and

$$N_3(a, b) = \frac{a + \sqrt{ab} + b}{3}.$$

The expression $N_3(a, b)$ is famous as Heron's mean. The inequalities (10) admit many non-negative differences. Based on these differences the author [...] proved the following result:

$$D_{SA} \leq \begin{cases} \frac{1}{3}D_{SH} \leq \begin{cases} \frac{1}{2}D_{AH} \leq \begin{cases} 4D_{N_2N_1} \leq \frac{4}{3}D_{N_2G} \leq D_{AG} \leq 4D_{AN_2} \\ \frac{1}{2}D_{SG} \leq D_{AG} \end{cases} \\ \frac{2}{3}D_{SN_1} \leq \frac{1}{2}D_{SG} \end{cases} \\ \frac{4}{5}D_{SN_2} \leq 4D_{AN_2} \\ \frac{3}{4}D_{SN_3} \leq \frac{2}{3}D_{SN_1} \end{cases}, \quad (11)$$

where $D_{SA} = S - A$, $D_{SH} = S - H$, etc. Some applications of the inequalities (11) can be seen in [6], [7].

Combining (9) and (10), we have the following sequence of inequalities:

$$P_1 \leq P_2 \leq P_3 \leq H \leq P_4 \leq G \leq N_1 \leq N_3 \leq N_2 \leq A \leq (P_5 \text{ or } S) \leq P_6. \quad (12)$$

The inequalities (12) admits many nonnegative differences such as

$$D_{tp}(a, b) = bg_{tp}\left(\frac{a}{b}\right) = b\left[f_t\left(\frac{a}{b}\right) - f_p\left(\frac{a}{b}\right)\right], \quad (13)$$

where

$$g_{tp}(x) = f_t(x) - f_p(x), \quad f_t(x) \geq f_p(x), \quad \forall x > 0.$$

More precisely, the expression (12) and the function $f : (0, \infty) \rightarrow \mathbb{R}$ given in (13) lead us to the following result:

$$\begin{aligned} f_{P_1}(x) &\leq f_{P_2}(x) \leq f_{P_3}(x) \leq f_H(x) \leq f_{P_4}(x) \leq f_G(x) \leq f_{N_1}(x) \\ &\leq f_{N_3}(x) \leq f_{N_2}(x) \leq f_A(x) \leq (f_{P_5}(x) \text{ or } f_S(x)) \leq f_{P_6}(x). \end{aligned} \quad (14)$$

Equivalently,

$$\begin{aligned} \frac{x(x^2+1)}{x^3+1} &\leq \frac{x(x+1)}{x^2+1} \leq \frac{x(\sqrt{x}+1)}{x^{3/2}+1} \leq \frac{2x}{1+x} \leq \frac{4x}{(\sqrt{x}+1)^2} \leq \\ &\leq \sqrt{x} \leq \left(\frac{\sqrt{x}+1}{2}\right)^2 \leq \frac{x+\sqrt{x}+1}{3} \leq \left(\frac{\sqrt{x}+1}{2}\right) \left(\sqrt{\frac{x+1}{2}}\right) \leq \\ &\leq \frac{x+1}{2} \leq \left(\left(\frac{x+1}{\sqrt{x}+1}\right)^2 \text{ or } \sqrt{\frac{x^2+1}{2}}\right) \leq \frac{x^2+1}{x+1}. \end{aligned} \quad (15)$$

In this paper our aim is to produce new inequalities for the difference of means arising due to inequalities given in (12). In another words we shall improve considerably the inequalities given in (11). For this we need first to know the convexity of the difference of means. In total, we have 77 differences. Some of them are equal to each other with some multiplicative constants. Some of them are not convex and some of them are convex.

2 Convexity of Difference of Means

Let us prove now the convexity of some of the difference of means arising due to inequalities (11). In order to prove it we shall make use of the following lemma (ref. Taneja [...]).

Lemma 2.1. *Let $f : I \subset \mathbb{R}_+ \rightarrow \mathbb{R}$ be a convex and differentiable function satisfying $f(1) = 0$. Consider a function*

$$\phi_f(a, b) = af\left(\frac{b}{a}\right), \quad a, b > 0,$$

then the function $\phi_f(a, b)$ is convex in $\mathbb{R}_+^2$. Additionally, if $f'(1) = 0$, then the following inequality hold:

$$0 \leq \phi_f(a, b) \leq \left(\frac{b-a}{a}\right) \phi'_f(a, b).$$

Inequalities appearing in (12) admits 77 nonnegative differences. Within these we have the following two set of equalities with multiplicative constants:

$$(i) \quad D_{P_6A} = D_{P_6H} = \frac{1}{2}D_{AH}.$$

$$(ii) \quad D_{P_5P_4} = D_{AN_3} = \frac{2}{3}D_{AN_1} = \frac{1}{3}D_{AG} = 2D_{N_3N_1} = \frac{1}{2}D_{N_3G} = \frac{2}{3}D_{N_1G}.$$

Not all the differences of means appearing in (12) are convex. We shall consider only those are convex. It is easy to check that in all the cases, $f_{(\cdot)}(1) = 1$, i.e., $g_{tp}(1) = 0$, $t > p$. According to Lemma 2.1, it is sufficient to show the convexity of the functions $g_{tp}(x)$, i.e., to show that the second order derivative of $g_{tp}(x)$ is nonnegative for all $x > 0$.

1. **For $D_{P_6S}(a, b)$:** We can write $D_{P_6S}(a, b) = b g_{P_6S}(a/b)$, where

$$g_{P_6S}(x) = f_{P_6}(x) - f_S(x) = \frac{x^2 + 1}{x + 1} - \sqrt{\frac{x^2 + 1}{2}}.$$

This gives

$$g''_{P_6S}(x) = \frac{2 \left[2(2x^2 + 2)^{3/2} - (x + 1)^3 \right]}{(x + 1)^3 (2x^2 + 2)^{3/2}}. \quad (16)$$

Since, $S \geq A$, this implies that $S^3 \geq A^3$, i.e., $\left(\sqrt{\frac{x^2+1}{2}}\right)^3 - \left(\frac{x+1}{2}\right)^3 \geq 0$. This gives $2(2x^2 + 2)^{3/2} - (x + 1)^3 \geq 0$. Thus we have $g''_{P_6S}(x) \geq 0$ for all $x > 0$.

2. **For $D_{P_6N_2}(a, b)$:** We can write $D_{P_6N_2}(a, b) = b g_{P_6N_2}(a/b)$, where

$$g_{P_6N_2}(x) = f_{P_6}(x) - f_{N_2}(x) = \frac{4(x^2 + 1) - \sqrt{2x + 2}(\sqrt{x} + 1)(x + 1)}{4(x + 1)}.$$

This gives

$$g''_{P_6N_2}(x) = \frac{(x+1)^3 (x^{3/2} + 1) + 16x^{3/2} (2x+2)^{3/2}}{4x^{3/2}(x+1)^3(2x+2)^{3/2}} > 0, \forall x > 0. \quad (17)$$

3. **For $\mathbf{D_{P_6N_3}(a, b)}$:** We can write $D_{P_6N_3}(a, b) = b g_{P_6N_3}(a/b)$, where

$$g_{P_6N_3}(x) = f_{P_6}(x) - f_{N_3}(x) = \frac{2(x^2 + 1) - \sqrt{x}(\sqrt{x} + 1)^2}{3(x+1)}.$$

This gives

$$g''_{P_6N_3}(x) = \frac{48x^{3/2} + (x+1)^3}{12x^{3/2}(x+1)^3} > 0, \forall x > 0. \quad (18)$$

4. **For $\mathbf{D_{P_6N_1}(a, b)}$:** We can write $D_{P_6N_1}(a, b) = b g_{P_6N_1}(a/b)$, where

$$g_{P_6N_1}(x) = f_{P_6}(x) - f_{N_1}(x) = \frac{3(x^2 + 1) - 2\sqrt{x}(x + \sqrt{x} + 1)}{4(x+1)}.$$

This gives

$$g''_{P_6N_1}(x) = \frac{32x^{3/2} + (x+1)^3}{8x^{3/2}(x+1)^3} > 0, \forall x > 0. \quad (19)$$

5. **For $\mathbf{D_{P_6G}(a, b)}$:** We can write $D_{P_6G}(a, b) = b g_{P_6G}(a/b)$, where

$$\begin{aligned} g_{P_6G}(x) &= f_{P_6}(x) - f_G(x) = \frac{x^2 + 1}{x+1} - \sqrt{x} \\ &= \frac{(\sqrt{x} - 1)^2 (x + \sqrt{x} + 1)}{x+1}. \end{aligned}$$

This gives

$$g''_{P_6G}(x) = \frac{16x^{3/2} + (x+1)^3}{4x^{3/2}(x+1)^3} > 0, \forall x > 0. \quad (20)$$

6. **For $\mathbf{D_{P_6P_4}(a, b)}$:** We can write $D_{P_6P_4}(a, b) = b g_{P_6P_4}(a/b)$, where

$$\begin{aligned} g_{P_6P_4}(x) &= f_{P_6}(x) - f_{P_4}(x) = \frac{x^2 + 1}{x+1} - \frac{4x}{(\sqrt{x} + 1)^2} \\ &= \frac{(x+1)(x^2 - 4x + 1) + 2\sqrt{x}(x^2 + 1)}{(x+1)(\sqrt{x} + 1)^2}. \end{aligned}$$

This gives

$$g''_{P_6P_4}(x) = \frac{2[3(x^3 + 1) + 17x(x+1) + 2\sqrt{x}(x^2 + 6x + 1)]}{\sqrt{x}(\sqrt{x} + 1)^4(x+1)^3} > 0, \forall x > 0. \quad (21)$$

7. **For $\mathbf{D}_{P_6P_2}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_6P_2}(a, b) = b g_{P_6P_2}(a/b)$, where

$$g_{P_6P_2}(x) = f_{P_6}(x) - f_{P_2}(x) = \frac{(x-1)^2(x^2+x+1)}{(x+1)(x^2+1)}.$$

This gives

$$g''_{P_6P_2}(x) = \frac{2(x^6 + 15x^4 + 16x^3 + 15x^2 + 1)}{(x+1)^3(x^2+1)^3} > 0, \forall x > 0. \quad (22)$$

8. **For $\mathbf{D}_{P_6P_1}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_6P_1}(a, b) = b g_{P_6P_1}(a/b)$, where

$$g_{P_6P_1}(x) = f_{P_6}(x) - f_{P_1}(x) = \frac{(x-1)^2(x^2+1)}{(x+1)^3}.$$

This gives

$$g''_{P_6P_1}(x) = \frac{2(x^2+1) \left(\frac{((x-1)^2-x)^2+x^2+1}{(x-1)^4+x(x-1)^2} \right) + 8x^3}{(x^3+1)^3} > 0, \forall x > 0. \quad (23)$$

9. **For $\mathbf{D}_{P_5A}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5A}(a, b) = b g_{P_5A}(a/b)$, where

$$\begin{aligned} g_{P_5A}(x) &= f_{P_5}(x) - f_A(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \frac{x+1}{2} \\ &= \frac{(x+1)(\sqrt{x}-1)^2}{2(\sqrt{x}+1)^2}. \end{aligned}$$

This gives

$$g''_{P_5A}(x) = \frac{4\sqrt{x} \left[(\sqrt{x}-1)^2 + \sqrt{x} \right] + (x-1)^2}{2x^{3/2}(\sqrt{x}+1)^4} > 0, \forall x > 0. \quad (24)$$

10. **For $\mathbf{D}_{P_5N_2}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5N_2}(a, b) = b g_{P_5N_2}(a/b)$, where

$$g_{P_5N_2}(x) = f_{P_5}(x) - f_{N_2}(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \left(\frac{\sqrt{x}+1}{2} \right) \left(\sqrt{\frac{x+1}{2}} \right).$$

This gives

$$\begin{aligned} g''_{P_5N_2}(x) &= \frac{1}{8(\sqrt{x}+1)x^{3/2}(x+1)\sqrt{2x+2}} \times \\ &\times \left(\frac{(\sqrt{x}+1)^4(x^{3/2}+1) + 4\sqrt{2x+2}(x+1)}{\times \left[4\sqrt{x} \left((\sqrt{x}-1)^2 + \sqrt{x} \right) + (x-1)^2 \right]} \right) > 0, \forall x > 0. \quad (25) \end{aligned}$$

11. **For $\mathbf{D}_{P_5N_3}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5N_3}(a, b) = b g_{P_5N_3}(a/b)$, where

$$\begin{aligned} g_{P_5N_3}(x) &= f_{P_5}(x) - f_{N_3}(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \frac{x + \sqrt{x} + 1}{3} \\ &= \frac{(\sqrt{x}-1)^2 (2x + \sqrt{x} + 2)}{2(\sqrt{x}+1)^2}. \end{aligned}$$

This gives

$$g''_{P_5N_3}(x) = \frac{7(\sqrt{x}-1)^2 (x + 6\sqrt{x} + 1) + 40x}{12x^{3/2} (\sqrt{x}+1)^4} > 0, \forall x > 0. \quad (26)$$

12. **For $\mathbf{D}_{P_5N_1}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5N_1}(a, b) = b g_{P_5N_1}(a/b)$, where

$$\begin{aligned} g_{P_5N_1}(x) &= f_{P_5}(x) - f_{N_1}(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \left(\frac{\sqrt{x}+1}{2} \right)^2 \\ &= \frac{(\sqrt{x}-1)^2 (3x + 2\sqrt{x} + 3)}{4(\sqrt{x}+1)^2}. \end{aligned}$$

This gives

$$g''_{P_5N_1}(x) = \frac{5(\sqrt{x}-1)^2 (x + 6\sqrt{x} + 1) + 32x}{8x^{3/2} (\sqrt{x}+1)^4} > 0, \forall x > 0. \quad (27)$$

13. **For $\mathbf{D}_{P_5G}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5G}(a, b) = b g_{P_5G}(a/b)$, where

$$\begin{aligned} g_{P_5G}(x) &= f_{P_5}(x) - f_G(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \sqrt{x} \\ &= \frac{(\sqrt{x}-1)^2 (x + \sqrt{x} + 1)}{(\sqrt{x}+1)^2}. \end{aligned}$$

This gives

$$g''_{P_5G}(x) = \frac{3[4\sqrt{x}(x+1) + (x-1)^2]}{4x^{3/2} (\sqrt{x}+1)^4} > 0, \forall x > 0. \quad (28)$$

14. **For $\mathbf{D}_{P_5H}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5H}(a, b) = b g_{P_5H}(a/b)$, where

$$g_{P_5H}(x) = f_{P_5}(x) - f_H(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \frac{2x}{x+1}.$$

This gives

$$\begin{aligned} g''_{P_5H}(x) &= \frac{1}{2x^{3/2} (x+1)^3 (\sqrt{x}+1)^4} \times \\ &\times \left((x+1) [(x-1)^4 + 16x^2] + 4\sqrt{x} [(x+1)^4 + 2x(x^2 + 6x + 1)] \right) > 0, \forall x > 0. \end{aligned} \quad (29)$$

15. **For $\mathbf{D}_{P_5P_3}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5P_3}(a, b) = b g_{P_5P_3} \left(\frac{a}{b} \right)$, where

$$\begin{aligned} g_{P_5P_3} &= f_{P_5}(x) - f_{P_3}(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \frac{x(\sqrt{x}+1)}{x^{3/2}+1} \\ &= \frac{(\sqrt{x}-1)^2 (x^2 + x^{3/2} + 3x + \sqrt{x} + 1)}{(\sqrt{x}+1)(x^{3/2}+1)}. \end{aligned}$$

This gives

$$g''_{P_5P_3}(x) = \frac{s_1(x)}{4x^{3/2} (x^{3/2} + 1)^3 (\sqrt{x} + 1)}, \quad (30)$$

where

$$s_1(x) = \left(\begin{array}{l} 2x^5 + 2x^{9/2} - 27x^4 + 75x^{7/2} - 123x^3 + \\ + 198x^{5/2} - 123x^2 + 75x^{3/2} - 27x + 2\sqrt{x} + 2 \end{array} \right).$$

Now we shall show that $s_1(x) > 0$, $\forall x > 0$. Let us consider

$$h_1(t) = s_1(t^2) = \left(\begin{array}{l} 2t^{10} + 2t^9 - 27t^8 + 75t^7 - 123t^6 + \\ + 198t^5 - 123t^4 + 75t^3 - 27t^2 + 2t + 2 \end{array} \right).$$

The polynomial equation $h(t) = 0$ of 10^{th} degree admits 10 solutions. Out of them 8 are complex and are given by

$$\begin{aligned} &0.002340265888 \pm 0.6477415967 I; \quad 0.005577710358 \pm 1.543805357 I; \\ &0.4030868619 \pm 0.2323740013 I; \quad 1.862033520 \pm 1.073436573 I. \end{aligned}$$

The real solutions are -5.359491864 and -0.1865848527 . Both these solutions are negative. Since we are working with $t > 0$, this means that there are no real positive solutions of the equation $h_1(t) = 0$. Thus we conclude that either $h_1(t) > 0$ or $h_1(t) < 0$, for all $t > 0$. In order to check it is sufficient to see for any particular value of $h_1(t)$, for example when $t = 1$. This gives $h_1(1) = 56$, hereby proving that $h_1(t) > 0$ for all $t > 0$, consequently, $s_1(x) > 0$, for all $x > 0$. Finally, we have $g''_{P_5P_3}(x) > 0$, $\forall x > 0$.

16. **For $\mathbf{D}_{P_5P_2}(\mathbf{a}, \mathbf{b})$:** We can write $D_{P_5P_2}(a, b) = b g_{P_5P_2} \left(\frac{a}{b} \right)$, where

$$\begin{aligned} g_{P_5P_2} &= f_{P_5}(x) - f_{P_2}(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \frac{x(x+1)}{x^2+1} \\ &= \frac{(x+1)(x^{3/2}-1)^2}{(\sqrt{x}+1)^2(x^2+1)}. \end{aligned}$$

This gives

$$g''_{P_5P_2}(x) = \frac{s_2(x)}{2x^{3/2} (x^2 + 1)^3 (\sqrt{x} + 1)^4}, \quad (31)$$

where

$$s_2(x) = \left(\begin{array}{l} x^8 + 4x^{15/2} - 6x^7 - 12x^6 + 14x^5 + 92x^{9/2} + \\ + 102x^4 + 92x^{7/2} + 14x^3 - 12x^2 - 6x + 4\sqrt{x} + 1 \end{array} \right).$$

Now we shall show that $s_2(x) > 0$, $\forall x > 0$. Let us consider

$$h_2(t) = s_2(t^2) = \left(\begin{array}{l} t^{16} + 4t^{15} - 6t^{14} - 12t^{12} + 14t^{10} + 92t^9 + \\ + 102t^8 + 92t^7 + 14t^6 - 12t^4 - 6t^2 + 4t + 1 \end{array} \right).$$

The polynomial equation $h_2(t) = 0$ of 16^{th} degree admits 16 solutions. Out of them 14 are complex (not written here) and two of them are real given by -5.230974171 and -0.1911689806 . Both these solutions are negative. Since we are working with $t > 0$, this means that there are no real positive solutions of the equation $h_2(t) = 0$. Thus we conclude that either $h_2(t) > 0$ or $h_2(t) < 0$, for all $t > 0$. In order to check it is sufficient to see for any particular value of $h_2(t)$, for example when $t = 1$. This gives $h_2(1) = 288$, hereby proving that $h_2(t) > 0$ for all $t > 0$, consequently, $s_2(x) > 0$, for all $x > 0$. Finally, we have $g''_{P_5P_2}(x) > 0$, $\forall x > 0$.

17. **For $D_{P_5P_1}(a, b)$:** We can write $D_{P_5P_1}(a, b) = b g_{P_5P_1}(\frac{a}{b})$, where

$$\begin{aligned} g_{P_5P_1} &= f_{P_5}(x) - f_{P_1}(x) = \left(\frac{x+1}{\sqrt{x}+1} \right)^2 - \frac{x(x^2+1)}{x^3+1} \\ &= \frac{\left(\begin{array}{l} x^4 + 2x^{7/2} + 4x^3 + 4x^{5/2} + \\ + 4x^2 + 4x^{3/2} + 4x + 2\sqrt{x} + 1 \end{array} \right) (\sqrt{x}-1)^2}{(\sqrt{x}+1)^2 (x^{3/2}+1)}. \end{aligned}$$

This gives

$$g''_{P_5P_1}(x) = \frac{s_3(x)}{2x^{3/2} (x^3+1)^3 (\sqrt{x}+1)^4}, \quad (32)$$

where

$$s_3(x) = \left(\begin{array}{l} x^{11} + 4x^{21/2} - 6x^{10} + 4x^{19/2} + x^9 - 12x^{17/2} - \\ - 45x^8 - 36x^{15/2} + 30x^7 + 144x^{13/2} + 99x^6 + \\ + 48x^{11/2} + 99x^5 + 144x^{9/2} + 30x^4 - 36x^{7/2} - \\ - 45x^3 - 12x^{5/2} + x^2 + 4x^{3/2} - 6x + 4\sqrt{x} + 1 \end{array} \right).$$

Now we shall show that $s_3(x) > 0$, $\forall x > 0$. Let us consider

$$h_3(t) = s_3(t^2) = \left(\begin{array}{l} t^{22} + 4t^{21} - 6t^{20} + 4t^{19} + t^{18} - 12t^{17} - \\ - 45t^{16} - 36t^{15} + 30t^{14} + 144t^{13} + 99t^{12} + \\ + 48t^{11} + 99t^{10} + 144t^9 + 30t^8 - 36t^7 - \\ - 45t^6 - 12t^5 + t^4 + 4t^3 - 6t^2 + 4t + 1 \end{array} \right).$$

The polynomial equation $h_3(t) = 0$ of 22^{nd} degree admits 22 solutions. Out of them 20 are complex (not written here) and two of them are real given by -5.269597986 and -0.1897677968 . Both these solutions are negative. Since we are working with $t > 0$, this means that there are no real positive solutions of the equation $h_3(t) = 0$. Thus we conclude that either $h_3(t) > 0$ or $h_3(t) < 0$, for all $t > 0$. In order to check it is sufficient to see for any particular value of $h_3(t)$, for example when $t = 1$. This gives $h_3(1) = 416$, hereby proving that $h_3(t) > 0$ for all $t > 0$, consequently, $s_3(x) > 0$, for all $x > 0$. Finally, we have $g''_{P_5P_1}(x) > 0, \forall x > 0$.

18. **For $D_{SP_4}(\mathbf{a}, \mathbf{b})$:** We can write $D_{SP_4}(a, b) = b g_{SP_4}(\frac{a}{b})$, where

$$g_{SP_4} = f_S(x) - f_{P_4}(x) = \sqrt{\frac{x^2 + 1}{2}} - \frac{4x}{(\sqrt{x} + 1)^2}.$$

This gives

$$g''_{SP_4}(x) = \frac{2 \left[\sqrt{x} (\sqrt{x} + 1)^4 + 3 (2x^2 + 2)^{3/2} \right]}{\sqrt{x} (\sqrt{x} + 1)^4 (2x^2 + 2)^{3/2}} > 0, \forall x > 0. \quad (33)$$

19. **For $D_{AP_4}(\mathbf{a}, \mathbf{b})$:** We can write $D_{AP_4}(a, b) = b g_{AP_4}(a/b)$, where

$$g_{AP_4}(x) = f_A(x) - f_{P_4}(x) = \frac{x + 1}{2} - \frac{4x}{(\sqrt{x} + 1)^2}.$$

This gives

$$g''_{AP_4}(x) = \frac{6}{\sqrt{x} (\sqrt{x} + 1)^4} > 0, \forall x > 0. \quad (34)$$

20. **For $D_{SA}(\mathbf{a}, \mathbf{b})$:** We can write $D_{SA}(a, b) = b g_{SA}(a/b)$, where

$$g_{SA}(x) = f_S(x) - f_A(x) = \sqrt{\frac{x^2 + 1}{2}} - \frac{x + 1}{2}.$$

This gives

$$g''_{SA}(x) = \frac{1}{(x^2 + 1) \sqrt{2x^2 + 2}} > 0, \forall x > 0. \quad (35)$$

21. **For $D_{SN_2}(\mathbf{a}, \mathbf{b})$:** We can write $D_{SN_2}(a, b) = b g_{SN_2}(a/b)$, where

$$g_{SN_2}(x) = f_S(x) - f_{N_2}(x) = \sqrt{\frac{x^2 + 1}{2}} - \sqrt{\frac{x + 1}{2}} \left(\frac{\sqrt{x} + 1}{2} \right).$$

This gives

$$g''_{SN_2}(x) = \frac{1}{8x^{3/2} (x^2 + 1) (x + 1) \sqrt{2x^2 + 2} \sqrt{2x + 2}} \times \left(\frac{\sqrt{2x^2 + 2} (x^{3/2} + 1) (x^2 + 1) +}{+ 8x^{3/2} (x + 1) \sqrt{2x + 2}} \right), \forall x > 0. \quad (36)$$

22. **For $D_{SN_1}(\mathbf{a}, \mathbf{b})$:** We can write $D_{SN_1}(a, b) = b g_{SN_1}(a/b)$, where

$$g_{SN_1}(x) = f_S(x) - f_{N_1}(x) = \sqrt{\frac{x^2 + 1}{2}} - \left(\frac{\sqrt{x} + 1}{2}\right)^2.$$

This gives

$$g''_{SN_1}(x) = \frac{8x^{3/2} + (x^2 + 1)\sqrt{2x^2 + 2}}{8x^{3/2}(x^2 + 1)\sqrt{2x^2 + 2}} > 0, \forall x > 0. \quad (37)$$

23. **For $D_{AN_2}(\mathbf{a}, \mathbf{b})$:** We can write $D_{AN_2}(a, b) = b g_{AN_2}(a/b)$, where

$$g_{AN_2}(x) = f_A(x) - f_{N_2}(x) = \frac{x + 1}{2} - \sqrt{\frac{x + 1}{2}} \left(\frac{\sqrt{x} + 1}{2}\right).$$

This gives

$$g''_{AN_2}(x) = \frac{x^{3/2} + 1}{8x^{3/2}(x + 1)\sqrt{2x + 2}} > 0, \forall x > 0. \quad (38)$$

24. **For $D_{AG}(\mathbf{a}, \mathbf{b})$:** We can write $D_{AG}(a, b) = b g_{AG}(a/b)$, where

$$g_{AG}(x) = f_A(x) - f_G(x) = \frac{x + 1}{2} - \sqrt{x} = \frac{(\sqrt{x} - 1)^2}{2}.$$

This gives

$$g''_{AG}(x) = \frac{1}{4x^{3/2}} > 0, \forall x > 0. \quad (39)$$

25. **For $D_{AH}(\mathbf{a}, \mathbf{b})$:** We can write $D_{AH}(a, b) = b g_{AH}(a/b)$, where

$$g_{AH}(x) = f_A(x) - f_H(x) = \frac{x + 1}{2} - \frac{2x}{x + 1} = \frac{(x - 1)^2}{2(x + 1)}.$$

This gives

$$g''_{AH}(x) = \frac{4}{(x + 1)^3} > 0, \forall x > 0. \quad (40)$$

26. **For $D_{N_2N_1}(\mathbf{a}, \mathbf{b})$:** We can write $D_{N_2N_1}(a, b) = b g_{N_2N_1}(a/b)$, where

$$g_{N_2N_1}(x) = f_{N_2}(x) - f_{N_1}(x) = \sqrt{\frac{x^2 + 1}{2}} - \left(\frac{\sqrt{x} + 1}{2}\right)^2.$$

This gives

$$g''_{N_2N_1}(x) = \frac{(x + 1)\sqrt{2x + 2} - (x^{3/2} + 1)}{8x^{3/2}(x + 1)\sqrt{2x + 2}}. \quad (41)$$

From (14), we have $(x + 1) \geq \left(\frac{\sqrt{x}+1}{2}\right) \sqrt{2x+2}$. This gives

$$\begin{aligned} & (x + 1)\sqrt{2x+2} - (x^{3/2} + 1) \\ &= (x + 1)\sqrt{2x+2} - (\sqrt{x} + 1)(x - \sqrt{x} + 1) \\ &\geq \left(\frac{\sqrt{x}+1}{2}\right) [2x+2 - 2(x - \sqrt{x} + 1)] \\ &\geq \sqrt{x}(\sqrt{x} + 1) > 0, \forall x > 0. \end{aligned}$$

This proves that $g''_{N_2 N_1}(x) > 0, \forall x > 0$.

27. **For $D_{SG}(a, b)$:** We can write $D_{SG}(a, b) = b g_{SG}(a/b)$, where

$$g_S(x) = f_S(x) - f_G(x) = \sqrt{\frac{x^2+1}{2}} - \sqrt{x}.$$

This gives

$$g''_{AH}(x) = \frac{4x^{3/2} + \sqrt{2x^2+2}(x^2+1)}{4x^{3/2}\sqrt{2x^2+2}(x^2+1)} > 0, \forall x > 0. \quad (42)$$

The convexity of the expressions given in parts 20-27 is already given in [11] but we have written them again because we need them in the next section.

3 Sequences of Inequalities

In this section we shall bring sequence of inequalities based on the differences arising due to (8). The results given in this section are based on the applications of the following lemma [11]:

Lemma 3.1. *Let $f_1, f_2 : I \subset \mathbb{R}_+ \rightarrow \mathbb{R}$ be two convex functions satisfying the assumptions:*

- (i) $f_1(1) = f'_1(1) = 0, f_2(1) = f'_2(1) = 0$;
- (ii) f_1 and f_2 are twice differentiable in $\mathbb{R}_+$;
- (iii) there exists the real constants α, β such that $0 \leq \alpha < \beta$ and

$$\alpha \leq \frac{f''_1(x)}{f''_2(x)} \leq \beta, f''_2(x) > 0,$$

for all $x > 0$ then we have the inequalities:

$$\alpha \phi_{f_2}(a, b) \leq \phi_{f_1}(a, b) \leq \beta \phi_{f_2}(a, b),$$

for all $a, b \in (0, \infty)$, where the function $\phi_{(\cdot)}(a, b)$ is as defined in Lemma 2.1.

Remark 3.1. From the above lemma we observe that

$$\eta(x) = \beta f_2(x) - f_1(x), \forall x > 0.$$

This gives $\eta(1) = \eta'(1) = 0$. In order to obtain β , let us consider $\eta''(1) = 0$. This gives $\beta = f_1''(1)/f_2''(1)$. We shall use this argument to prove the results given in the theorem below.

Theorem 3.1. The following sequences of inequalities hold:

$$\begin{aligned} \frac{1}{8}D_{P_6P_1} &\leq \frac{1}{6}D_{P_6P_2} \leq D_{SA} \leq \frac{1}{3}D_{SH} \leq \frac{1}{2}D_{AH} \leq \\ &\leq \left\{ \frac{4}{9}D_{P_6N_2} \right\} \leq \left\{ \frac{3}{7}D_{P_6N_3} \right\} \leq \left\{ \frac{2}{5}D_{P_6N_1} \right\} \leq \frac{1}{3}D_{P_6G} \leq \left\{ \frac{2}{3}D_{P_5H} \right\} \leq \\ &\leq 4D_{N_2N_1} \leq \frac{4}{3}D_{N_2G} \leq D_{AG} \leq 4D_{AN_2} \leq \frac{2}{3}D_{P_5G} \leq \\ &\leq D_{P_5N_1} \leq \frac{6}{5}D_{P_5N_3} \leq \frac{4}{3}D_{P_5N_2} \leq 2D_{P_5A}, \end{aligned} \quad (43)$$

$$D_{SA} \leq \left\{ \frac{4}{3}D_{SN_2} \right\} \leq \frac{2}{3}D_{SN_1} \leq \left\{ \frac{1}{3}D_{P_6G} \right\} \leq \frac{2}{5}D_{P_5H} \quad (44)$$

and

$$\left\{ \frac{1}{8}D_{P_6P_1} \right\} \leq \left\{ \frac{1}{6}D_{P_6P_2} \right\} \leq \frac{2}{7}D_{P_5P_3} \leq \frac{4}{9}D_{P_6N_2} \leq D_{P_6S} \leq D_{AG}. \quad (45)$$

Proof. We shall prove each part separately.

1. **For $D_{P_6P_1} \leq \frac{4}{3}D_{P_6P_2}$:** Let us consider the function $g_{P_6P_1-P_6P_2}(x) = f_{P_6P_1}''(x)/f_{P_6P_2}''(x)$, where $f_{P_6P_1}''(x)$ and $f_{P_6P_2}''(x)$ are as given by (23) and (22) respectively. This gives

$$\beta_{P_6P_1-P_6P_2} = g_{P_6P_1-P_6P_2}(1) = \frac{f_{P_6P_1}''(1)}{f_{P_6P_2}''(1)} = \frac{4}{3}.$$

In order to prove this part we shall show that $\frac{4}{3}D_{P_6P_2} - D_{P_6P_1} \geq 0$. We can write

$$\frac{4}{3}D_{P_6P_2} - D_{P_6P_1} = bf_{P_6P_2-P_6P_1}\left(\frac{a}{b}\right),$$

where

$$f_{P_6P_2-P_6P_1}(x) = \frac{4}{3}f_{P_6P_2}(x) - f_{P_6P_1}(x) = \frac{(x+1)^2(x-1)^4}{(x^3+1)(x^2+1)}.$$

Since $f_{P_6P_2-P_6P_1}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

2. **For $\mathbf{D}_{P_6P_2} \leq 6\mathbf{D}_{SA}$:** By considering the function $g_{P_6P_2SA}(x) = f''_{P_6P_2}(x)/f''_{SA}(x)$, where $f''_{P_6P_2}(x)$ and $f''_{SA}(x)$ are as given by (22) and (35) respectively, we get $\beta_{P_6P_1SA} = g_{P_6P_1SA}(1) = 6$. We can write

$$6D_{SA} - D_{P_6P_2} = b f_{SA-P_6P_2} \left(\frac{a}{b} \right),$$

where

$$f_{SA-P_6P_2}(x) = 6f_{SA}(x) - f_{P_6P_2}(x) = \frac{k_2(x)}{(x^2 + 1)(x + 1)},$$

with

$$k_2(x) = 3\sqrt{2x+2}(x+1)(x^2+1) - (4x^4 + 5x^3 + 6x^2 + 5x + 4).$$

Now we shall show that $k_2(x) > 0, \forall x > 0, x \neq 1$. Let us consider

$$h_2(x) = \left[3\sqrt{2x+2}(x+1)(x^2+1) \right]^2 - (4x^4 + 5x^3 + 6x^2 + 5x + 4)^2.$$

After simplifications we have

$$h_2(x) = (2x^4 + 4x^3 + 3x^2 + 4x + 2)(x-1)^4.$$

Since $h_2(x) > 0$ giving $k_2(x) > 0, \forall x > 0, x \neq 1$. Thus we have $f_{P_6P_2-P_6P_1}(x) > 0, \forall x > 0, x \neq 1$, thereby proving the required result.

Argument: Let a and b two positive numbers, i.e., $a > 0$ and $b > 0$. If $a^2 - b^2 > 0$, then we can conclude that $a > b$ because $a - b = (a^2 - b^2)/(a + b)$. We have used this argument to prove $k_2(x) > 0, \forall x > 0, x \neq 1$. We shall use frequently this argument to prove the other parts of the theorem.

3. **For $\mathbf{D}_{SA} \leq \frac{1}{3}\mathbf{D}_{SH}$:** This result is already appearing in (11).
 4. **For $\mathbf{D}_{SH} \leq \frac{3}{2}\mathbf{D}_{AH}$:** This result is already appearing in (11).
 5. **For $\mathbf{D}_{AH} \leq \frac{8}{9}\mathbf{D}_{P_6N_2}$:** By considering the function $g_{AH-P_6N_2}(x) = f''_{AH}(x)/f''_{P_6N_2}(x)$, we get $\beta_{AH-P_6N_2} = g_{AH-P_6N_2}(1) = \frac{8}{9}$, where $f''_{AH}(x)$ and $f''_{P_6N_2}(x)$ are as given by (40) and (17) respectively. We can write

$$\frac{8}{9}D_{P_6N_2} - D_{AH} = b f_{P_6N_2-AH} \left(\frac{a}{b} \right),$$

where

$$f_{P_6N_2-AH}(x) = \frac{8}{9}f_{P_6N_2}(x) - f_{AH}(x) = \frac{k_5(x)}{18(x+1)},$$

with

$$k_5(x) = 7x^2 + 18x + 7 - 4\sqrt{2x+2}(\sqrt{x}+1)(x+1).$$

Now we shall show that $k_5(x) > 0, \forall x > 0, x \neq 1$. Let us consider

$$h_5(x) = (7x^2 + 18x + 7)^2 - \left[4\sqrt{2x+2} (\sqrt{x} + 1) (x+1) \right]^2.$$

After simplifications we have

$$h_5(x) = (17x^2 + 4x^{3/2} + 38x + 4\sqrt{x} + 17) (\sqrt{x} - 1)^4.$$

Since $h_5(x) > 0$ giving $k_5(x) > 0, \forall x > 0, x \neq 1$. Thus we have $f_{P_6P_2-AH}(x) > 0, \forall x > 0, x \neq 1$, thereby proving the required result.

6. **For $D_{AH} \leq \frac{6}{7}D_{P_6N_3}$:** By considering the function $g_{AH-P_6N_3}(x) = f''_{AH}(x)/f''_{P_6N_3}(x)$, we get $\beta_{AH-P_6N_3} = g_{AH-P_6N_3}(1) = \frac{6}{7}$, where $f''_{AH}(x)$ and $f''_{P_6N_3}(x)$ are as given by (40) and (18) respectively. We can write

$$\frac{6}{7}D_{P_6N_3} - D_{AH} = b f_{P_6N_3-AH} \left(\frac{a}{b} \right),$$

where

$$f_{P_6N_3-AH}(x) = \frac{6}{7}f_{P_6N_3}(x) - f_{AH}(x) = \frac{(\sqrt{x}-1)^2}{18(x+1)}.$$

Since $f_{P_6N_3-AH}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

7. **For $D_{AH} \leq \frac{2}{3}D_{SP_4}$:** By considering the function $g_{AH-SP_4}(x) = f''_{AH}(x)/f''_{SP_4}(x)$, we get $\beta_{AH-SP_4} = g_{AH-SP_4}(1) = \frac{2}{3}$, where $f''_{AH}(x)$ and $f''_{SP_4}(x)$ are as given by (40) and (33) respectively. We can write

$$\frac{2}{3}D_{SP_4} - D_{AH} = b f_{SP_4-AH} \left(\frac{a}{b} \right),$$

where

$$f_{SP_4-AH}(x) = \frac{2}{3}f_{SP_4}(x) - f_{AH}(x) = \frac{k_7(x)}{10(x+1)(\sqrt{x}+1)^2},$$

with

$$k_7(x) = 4(x+1)\sqrt{2x+2}(\sqrt{x}+1)^2 - \left[5x^3 + 10x^{5/2} + 17x^2 + 17x + 10x(\sqrt{x}-1)^2 + 10\sqrt{x} + 5 \right].$$

Now we shall show that $k_7(x) > 0, \forall x > 0, x \neq 1$. Let us consider

$$h_7(x) = \left[4(x+1)\sqrt{2x+2}(\sqrt{x}+1)^2 \right]^2 - \left[5x^3 + 10x^{5/2} + 17x^2 + 17x + 10x(\sqrt{x}-1)^2 + 10\sqrt{x} + 5 \right]^2.$$

After simplifications we have

$$h_7(x) = (7x^3 + 70x^{5/2} + 201x^2 + 340x^{3/2} + 201x + 70\sqrt{x} + 7) (\sqrt{x} - 1)^6$$

Since $h_7(x) > 0$ giving $k_7(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{SP_4-AH}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

8. **For $\mathbf{D}_{P_6N_3} \leq \frac{14}{15}\mathbf{D}_{P_6N_1}$** : By considering the function $g_{P_6N_3-P_6N_1}(x) = f''_{P_6N_3}(x)/f''_{P_6N_1}(x)$, we get $\beta_{P_6N_3-P_6N_1} = g_{P_6N_3-P_6N_1}(1) = \frac{14}{15}$, where $f''_{P_6N_3}(x)$ and $f''_{P_6N_1}(x)$ are as given by (18) and (19) respectively. We can write

$$\frac{14}{15}D_{P_6N_1} - D_{P_6N_3} = b f_{P_6N_1-P_6N_3} \left(\frac{a}{b} \right),$$

where

$$f_{P_6N_1-P_6N_3}(x) = \frac{14}{15}f_{P_6N_1}(x) - f_{P_6N_3}(x) = \frac{(\sqrt{x} - 1)^2}{30(x + 1)}.$$

Since $f_{P_6N_1-P_6N_3}(x) > 0$, $\forall x > 0$, $x \neq 1$, hence proving the required result.

9. **For $\mathbf{D}_{P_6N_3} \leq \frac{2}{3}\mathbf{D}_{P_6P_4}$** : By considering the function $g_{P_6N_3-P_6P_4}(x) = f''_{P_6N_3}(x)/f''_{P_6P_4}(x)$, we get $\beta_{P_6N_3-P_6P_4} = g_{P_6N_3-P_6P_4}(1) = \frac{2}{3}$, where $f''_{P_6N_3}(x)$ and $f''_{P_6P_4}(x)$ are as given by (18) and (21) respectively. We can write

$$\frac{2}{3}D_{P_6P_4} - D_{P_6N_3} = b f_{P_6P_4-P_6N_3} \left(\frac{a}{b} \right),$$

where

$$f_{P_6P_4-P_6N_3}(x) = \frac{2}{3}f_{P_6P_4}(x) - f_{P_6N_3}(x) = \frac{\sqrt{x}(\sqrt{x} - 1)^4}{3(\sqrt{x} + 1)^2(x + 1)}.$$

Since $f_{P_6P_4-P_6N_3}(x) > 0$, $\forall x > 0$, $x \neq 1$, hence proving the required result.

10. **For $\mathbf{D}_{SP_4} \leq \frac{4}{5}\mathbf{D}_{P_6N_1}$** : By considering the function $g_{SP_4-P_6N_1}(x) = f''_{SP_4}(x)/f''_{P_6N_1}(x)$, we get $\beta_{SP_4-P_6N_1} = g_{SP_4-P_6N_1}(1) = \frac{4}{5}$, where $f''_{SP_4}(x)$ and $f''_{P_6N_1}(x)$ are as given by (33) and (19) respectively. We can write

$$\frac{4}{5}D_{P_6N_1} - D_{SP_4} = b f_{P_6N_1-SP_4} \left(\frac{a}{b} \right),$$

where

$$f_{P_6N_1-SP_4}(x) = \frac{4}{5}f_{P_6N_1}(x) - f_{SP_4}(x) = \frac{k_{10}(x)}{4(x + 1)(\sqrt{x} + 1)^2},$$

with

$$k_{10}(x) = 3x^3 + 4x^{5/2} + 9x^2 + 4x(\sqrt{x} - 1)^2 + 9x + 4\sqrt{x} + 3 - 2\sqrt{2x^2 + 2}(x + 1)(\sqrt{x} + 1)^2.$$

Now we shall show that $k_{10}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{10}(x) = \left[3x^3 + 4x^{5/2} + 9x^2 + 4x(\sqrt{x} - 1)^2 + 9x + 4\sqrt{x} + 3 \right]^2 - \left[2\sqrt{2x^2 + 2}(x + 1)(\sqrt{x} + 1)^2 \right]^2.$$

After simplifications we have

$$h_{10}(x) = (\sqrt{x} - 1)^4 \left(x^3(\sqrt{x} - 2)^2 + 4x^3 + 20x^{5/2} + 78x^2 + 20x^{3/2} + 4x + (2\sqrt{x} - 1)^2 \right).$$

Since $h_{10}(x) > 0$ giving $k_{10}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_6 N_1 - SP_4}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

11. **For $D_{SP_4} \leq \frac{5}{7}D_{P_6 P_4}$** : By considering the function $g_{SP_4 - P_6 P_4}(x) = f''_{SP_4}(x)/f''_{P_6 P_4}(x)$, we get $\beta_{SP_4 - P_6 P_4} = g_{SP_4 - P_6 P_4}(1) = \frac{5}{7}$, where $f''_{SP_4}(x)$ and $f''_{P_6 P_4}(x)$ are as given by (33) and (21) respectively. We can write

$$\frac{5}{7}D_{P_6 P_4} - D_{SP_4} = b f_{P_6 P_4 - SP_4} \left(\frac{a}{b} \right),$$

where

$$f_{P_6 P_4 - SP_4}(x) = \frac{5}{7}f_{P_6 P_4}(x) - f_{SP_4}(x) = \frac{k_{11}(x)}{14(x + 1)(\sqrt{x} + 1)^2},$$

with

$$k_{11}(x) = 10x^3 + 20x^{5/2} + 26x^2 + 26x + 20\sqrt{x} + 10 - 7\sqrt{2x^2 + 2}(\sqrt{x} + 1)^2(x + 1).$$

Now we shall show that $k_{11}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{11}(x) = (10x^3 + 20x^{5/2} + 26x^2 + 26x + 20\sqrt{x} + 10)^2 - \left[7\sqrt{2x^2 + 2}(\sqrt{x} + 1)^2(x + 1) \right]^2.$$

After simplifications we have

$$h_{11}(x) = 2(\sqrt{x} - 1)^4 \left(x^4 + 8x^{7/2} + 94x^3 + 264x^{5/2} + 386x^2 + 264x^{3/2} + 94x + 8\sqrt{x} + 1 \right).$$

Since $h_{11}(x) > 0$ giving $k_{11}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_6 P_4 - SP_4}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

12. **For $\mathbf{D}_{P_6N_2} \leq \frac{3}{4}\mathbf{D}_{P_6G}$** : By considering the function $g_{P_6N_2-P_6G}(x) = f''_{P_6N_2}(x)/f''_{P_6G}(x)$, we get $\beta_{P_6N_2-P_6G} = g_{P_6N_2-P_6G}(1) = \frac{3}{4}$, where $f''_{P_6N_2}(x)$ and $f''_{P_6G}(x)$ are as given by (17) and (20) respectively. We can write

$$\frac{3}{4}D_{P_6G} - D_{P_6N_2} = b f_{P_6G-P_6N_2} \left(\frac{a}{b} \right),$$

where

$$f_{P_6G-P_6N_2}(x) = \frac{3}{4}f_{P_6G}(x) - f_{P_6N_2}(x) = \frac{k_{12}(x)}{4(x+1)},$$

with

$$k_{12}(x) = \sqrt{2x+2}(\sqrt{x}+1)(x+1) - (x^2+1+3x^{3/2}+3\sqrt{x}).$$

Now we shall show that $k_5(x) > 0, \forall x > 0, x \neq 1$. Let us consider

$$h_{12}(x) = \left[\sqrt{2x+2}(\sqrt{x}+1)(x+1) \right]^2 - (x^2+1+3x^{3/2}+3\sqrt{x})^2.$$

After simplifications we have

$$h_{12}(x) = (\sqrt{x}-1)^4(x^2+2x^{3/2}+x+2\sqrt{x}+1).$$

Since $h_{12}(x) > 0$ giving $k_{12}(x) > 0, \forall x > 0, x \neq 1$. Thus we have $f_{P_6G-P_6N_2}(x) > 0, \forall x > 0, x \neq 1$, thereby proving the required result.

13. **For $\mathbf{D}_{P_6N_1} \leq \frac{5}{6}\mathbf{D}_{P_6G}$** : By considering the function $g_{P_6N_1-P_6G}(x) = f''_{P_6N_1}(x)/f''_{P_6G}(x)$, we get $\beta_{P_6N_1-P_6G} = g_{P_6N_1-P_6G}(1) = \frac{5}{6}$, where $f''_{P_6N_1}(x)$ and $f''_{P_6G}(x)$ are as given by (19) and (20) respectively. We can write

$$\frac{5}{6}D_{P_6G} - D_{P_6N_1} = b f_{P_6G-P_6N_1} \left(\frac{a}{b} \right),$$

where

$$f_{P_6G-P_6N_1}(x) = \frac{5}{6}f_{P_6G}(x) - f_{P_6N_1}(x) = \frac{(\sqrt{x}-1)^2}{12(x+1)}.$$

Since $f_{P_6G-P_6N_1}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

14. **For $\mathbf{D}_{P_6P_4} \leq \frac{7}{6}\mathbf{D}_{P_6G}$** : By considering the function $g_{P_6P_4-P_6G}(x) = f''_{P_6P_4}(x)/f''_{P_6G}(x)$, we get $\beta_{P_6P_4-P_6G} = g_{P_6P_4-P_6G}(1) = \frac{7}{6}$, where $f''_{P_6P_4}(x)$ and $f''_{P_6G}(x)$ are as given by (21) and (20) respectively. We can write

$$\frac{7}{6}D_{P_6G} - D_{P_6P_4} = b f_{P_6G-P_6P_4} \left(\frac{a}{b} \right),$$

where

$$f_{P_6G-P_6P_4}(x) = \frac{7}{6}f_{P_6G}(x) - f_{P_6P_4}(x) = \frac{(\sqrt{x}-1)^4[(\sqrt{x}-1)^2+\sqrt{x}]}{12(x+1)}.$$

Since $f_{P_6G-P_6P_4}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

15. **For $\mathbf{D}_{P_6G} \leq \frac{6}{5}\mathbf{D}_{P_5H}$:** By considering the function $g_{P_6G-P_5H}(x) = f''_{P_6G}(x)/f''_{P_5H}(x)$, we get $\beta_{P_6G-P_5H} = g_{P_6G-P_5H}(1) = \frac{6}{5}$, where $f''_{P_6G}(x)$ and $f''_{P_5H}(x)$ are as given by (20) and (29) respectively. We can write

$$\frac{6}{5}D_{P_5H} - D_{P_6G} = b f_{P_5H-P_6G}\left(\frac{a}{b}\right),$$

where

$$f_{P_5H-P_6G}(x) = \frac{6}{5}f_{P_5H}(x) - f_{P_6G}(x) = \frac{(\sqrt{x}-1)^4 \left[(\sqrt{x}-1)^2 + \sqrt{x} \right]}{5(x+1)}.$$

Since $f_{P_5H-P_6G}(x) > 0$, $\forall x > 0$, $x \neq 1$, hence proving the required result.

16. **For $\mathbf{D}_{P_6G} \leq 2\mathbf{D}_{AP_4}$:** By considering the function $g_{P_6G-AP_4}(x) = f''_{P_6G}(x)/f''_{AP_4}(x)$, we get $\beta_{P_6G-AP_4} = g_{P_6G-AP_4}(1) = 2$, where $f''_{P_6G}(x)$ and $f''_{AP_4}(x)$ are as given by (20) and (34) respectively. We can write

$$2D_{AP_4} - D_{P_6G} = b f_{AP_4-P_6G}\left(\frac{a}{b}\right),$$

where

$$f_{AP_4-P_6G}(x) = 2f_{AP_4}(x) - f_{P_6G}(x) = \frac{\sqrt{x}(\sqrt{x}-1)^4}{(\sqrt{x}+1)^2(x+1)}.$$

Since $f_{AP_4-P_6G}(x) > 0$, $\forall x > 0$, $x \neq 1$, hence proving the required result.

17. **For $\mathbf{D}_{P_5H} \leq 10\mathbf{D}_{N_2N_1}$:** By considering the function $g_{P_5H-N_2N_1}(x) = f''_{P_5H}(x)/f''_{N_2N_1}(x)$, we get $\beta_{P_5H-N_2N_1} = g_{P_5H-N_2N_1}(1) = 10$, where $f''_{P_5H}(x)$ and $f''_{N_2N_1}(x)$ are as given by (29) and (41) respectively. We can write

$$10D_{N_2N_1} - D_{P_5H} = b f_{N_2N_1-P_5H}\left(\frac{a}{b}\right),$$

where

$$f_{N_2N_1-P_5H}(x) = 10f_{N_2N_1}(x) - f_{P_5H}(x) = \frac{k_{17}(x)}{2(\sqrt{x}+1)^2(x+1)},$$

with

$$\begin{aligned} k_{17}(x) &= 5\sqrt{2x+2}(\sqrt{x}+1)^3(x+1) \\ &\quad - (7x^3 + 20x^{5/2} + 37x^2 + 32x^{3/2} + 37x + 20\sqrt{x} + 7). \end{aligned}$$

Now we shall show that $k_{17}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$\begin{aligned} h_{17}(x) &= \left[5\sqrt{2x+2}(\sqrt{x}+1)^3(x+1) \right]^2 \\ &\quad - (7x^3 + 20x^{5/2} + 37x^2 + 32x^{3/2} + 37x + 20\sqrt{x} + 7)^2. \end{aligned}$$

After simplifications we have

$$h_{17}(x) = (\sqrt{x} - 1)^4 \left(\begin{array}{l} x^4 + 24x^{7/2} + 72x^3 + 120x^{5/2} + \\ + 126x^2 + 120x^{3/2} + 72x + 24\sqrt{x} + 1 \end{array} \right).$$

Since $h_{17}(x) > 0$ giving $k_{17}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{N_2N_1-P_5H}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

18. **For $D_{AP_4} \leq 6D_{N_2N_1}$** : By considering the function $g_{AP_4-N_2N_1}(x) = f''_{AP_4}(x)/f''_{N_2N_1}(x)$, we get $\beta_{AP_4-N_2N_1} = g_{AP_4-N_2N_1}(1) = 6$, where $f''_{AP_4}(x)$ and $f''_{N_2N_1}(x)$ are as given by (34) and (41) respectively. We can write

$$6D_{N_2N_1} - D_{AP_4} = b f_{N_2N_1-AP_4} \left(\frac{a}{b} \right),$$

where

$$f_{N_2N_1-AP_4}(x) = 6f_{N_2N_1}(x) - f_{AP_4}(x) = \frac{k_{18}(x)}{2(\sqrt{x} + 1)^2},$$

with

$$k_{18}(x) = 3\sqrt{2x+2}(\sqrt{x}+1)^3 - 2(2x^2 + 7x^{3/2} + 6x + 7\sqrt{x} + 2).$$

Now we shall show that $k_{18}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{18}(x) = \left[3\sqrt{2x+2}(\sqrt{x}+1)^3 \right]^2 - (4x^2 + 14x^{3/2} + 12x + 14\sqrt{x} + 4)^2$$

After simplifications we have

$$h_{18}(x) = 2(\sqrt{x} - 1)^4 (x^2 + 2x^{3/2} + 2\sqrt{x} + 1).$$

Since $h_{18}(x) > 0$ giving $k_{18}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{N_2N_1-AP_4}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

19. **For $D_{N_2N_1} \leq \frac{1}{3}D_{N_2G}$** : This result is already appearing in (11).
 20. **For $D_{N_2G} \leq \frac{3}{4}D_{AG}$** : This result is already appearing in (11).
 21. **For $D_{AG} \leq 4D_{AN_2}$** : This result is already appearing in (11).
 22. **For $D_{AN_2} \leq \frac{1}{6}D_{P_5G}$** : By considering the function $g_{AN_2-P_5G}(x) = f''_{AN_2}(x)/f''_{P_5G}(x)$, we get $\beta_{AN_2-P_5G} = g_{AN_2-P_5G}(1) = \frac{1}{6}$, where $f''_{AN_2}(x)$ and $f''_{P_5G}(x)$ are as given by (38) and (28) respectively. We can write

$$\frac{3}{4}D_{P_5G} - D_{AN_2} = b f_{AN_2-P_5G} \left(\frac{a}{b} \right),$$

where

$$f_{AN_2-P_5G}(x) = \frac{1}{6}f_{P_5G}(x) - f_{AN_2}(x) = \frac{k_{22}(x)}{12(\sqrt{x}+1)},$$

with $k_{22}(x) = k_{18}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{N_2N_1-AP_4}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

23. **For $D_{P_5G} \leq \frac{3}{2}D_{P_5N_1}$:** By considering the function $g_{P_5G-P_5N_1}(x) = f_{P_5G}''(x)/f_{P_5N_1}''(x)$, we get $\beta_{P_5G-P_5N_1} = g_{P_5G-P_5N_1}(1) = \frac{3}{2}$, where $f_{P_5G}''(x)$ and $f_{P_5N_1}''(x)$ are as given by (28) and (27) respectively. We can write

$$\frac{3}{2}D_{P_5N_1} - D_{P_5G} = b f_{P_5G-P_5N_1}\left(\frac{a}{b}\right),$$

where

$$f_{P_5G-P_5N_1}(x) = \frac{3}{2}f_{P_5G}(x) - f_{P_5N_1}(x) = \frac{(\sqrt{x}-1)^2}{8(x+1)}.$$

Since $f_{P_5G-P_5N_1}(x) > 0$, $\forall x > 0$, $x \neq 1$, hence proving the required result.

24. **For $D_{P_5N_1} \leq \frac{6}{5}D_{P_5N_3}$:** By considering the function $g_{P_5N_1-P_5N_3}(x) = f_{P_5N_1}''(x)/f_{P_5N_3}''(x)$, we get $\beta_{P_5N_1-P_5N_3} = g_{P_5N_1-P_5N_3}(1) = \frac{6}{5}$, where $f_{P_5N_1}''(x)$ and $f_{P_5N_3}''(x)$ are as given by (27) and (26) respectively. We can write

$$\frac{6}{5}D_{P_5N_3} - D_{P_5N_1} = b f_{P_5N_3-P_5N_1}\left(\frac{a}{b}\right),$$

where

$$f_{P_5N_3-P_5N_1}(x) = \frac{6}{5}f_{P_5N_3}(x) - f_{P_5N_1}(x) = \frac{(\sqrt{x}-1)^2}{20(x+1)}.$$

Since $f_{P_5N_3-P_5N_1}(x) > 0$, $\forall x > 0$, $x \neq 1$, hence proving the required result.

25. **For $D_{P_5N_3} \leq \frac{10}{9}D_{P_5N_2}$:** By considering the function $g_{P_5N_3-P_5N_2}(x) = f_{P_5N_3}''(x)/f_{P_5N_2}''(x)$, we get $\beta_{P_5N_3-P_5N_2} = g_{P_5N_3-P_5N_2}(1) = \frac{10}{9}$, where $f_{P_5N_3}''(x)$ and $f_{P_5N_2}''(x)$ are as given by (26) and (25) respectively. We can write

$$\frac{10}{9}D_{P_5N_2} - D_{P_5N_3} = b f_{P_5N_2-P_5N_3}\left(\frac{a}{b}\right),$$

where

$$f_{P_5N_2-P_5N_3}(x) = \frac{10}{9}f_{P_5N_2}(x) - f_{P_5N_3}(x) = \frac{k_{25}(x)}{18(\sqrt{x}+1)^2},$$

with

$$k_{25}(x) = 2(4x^2 + 9x^{3/2} + 14x + 9\sqrt{x} + 4) - 5\sqrt{2x+2}(\sqrt{x}+1)^3.$$

Now we shall show that $k_{25}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{25}(x) = \left[2 \left(4x^2 + 9x^{3/2} + 14x + 9\sqrt{x} + 4 \right) \right]^2 - \left[5\sqrt{2x+2} (\sqrt{x} + 1)^3 \right]^2.$$

After simplifications we have

$$h_{25}(x) = 2 (\sqrt{x} - 1)^4 (7x^2 + 22x^{3/2} + 32x + 22\sqrt{x} + 7).$$

Since $h_{25}(x) > 0$ giving $k_{25}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_5N_2 \dashv P_5N_3}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

26. **For $\mathbf{D}_{P_5N_2} \leq \frac{3}{2}\mathbf{D}_{P_5A}$** : By considering the function $g_{P_5N_2 \dashv P_5A}(x) = f''_{P_5N_2}(x)/f''_{P_5A}(x)$, we get $\beta_{P_5N_2 \dashv P_5A} = g_{P_5N_2 \dashv P_5A}(1) = \frac{3}{2}$, where $f''_{P_5N_2}(x)$ and $f''_{P_5A}(x)$ are as given by (25) and (24) respectively. We can write

$$\frac{3}{2}D_{P_5A} - D_{P_5N_2} = b f_{P_5A \dashv P_5N_2} \left(\frac{a}{b} \right),$$

where

$$f_{P_5A \dashv P_5N_2}(x) = \frac{3}{2}f_{P_5A}(x) - f_{P_5N_2}(x) = \frac{k_{26}(x)}{4(\sqrt{x} + 1)^2},$$

with

$$k_{26}(x) = \sqrt{2x+2} (\sqrt{x} + 1)^3 - (x+1)(x+6\sqrt{x}+1).$$

Let us consider

$$h_{26}(x) = \left[\sqrt{2x+2} (\sqrt{x} + 1)^3 \right]^2 - \left[(x+1)(x+6\sqrt{x}+1) \right]^2.$$

After simplifications we have

$$h_{26}(x) = (\sqrt{x} - 1)^4 (x+1)(x+4\sqrt{x}+1).$$

Since $h_{26}(x) > 0$ giving $k_{26}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_5A \dashv P_5N_2}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

Parts 1-26 prove the sequences of inequalities appearing in (43).

27. **For $\mathbf{D}_{SA} \leq \frac{4}{5}\mathbf{D}_{SN_2}$** : This result is already appearing in (11).
 28. **For $\mathbf{D}_{SA} \leq \frac{3}{4}\mathbf{D}_{SN_3}$** : This result is already appearing in (11).

29. **For $\mathbf{D}_{\mathbf{SN}_2} \leq \frac{5}{6}\mathbf{D}_{\mathbf{SN}_1}$** : By considering the function $g_{\mathbf{SN}_2\text{-}\mathbf{SN}_1}(x) = f''_{\mathbf{SN}_2}(x)/f''_{\mathbf{SN}_1}(x)$, we get $\beta_{\mathbf{SN}_2\text{-}\mathbf{SN}_1} = g_{\mathbf{SN}_2\text{-}\mathbf{SN}_1}(1) = \frac{5}{6}$, where $f''_{\mathbf{SN}_2}(x)$ and $f''_{\mathbf{SN}_1}(x)$ are as given by (36) and (37) respectively. We can write

$$\frac{5}{6}D_{\mathbf{SN}_1} - D_{\mathbf{SN}_2} = b f_{\mathbf{SN}_1\text{-}\mathbf{SN}_2} \left(\frac{a}{b} \right),$$

where

$$f_{\mathbf{SN}_1\text{-}\mathbf{SN}_2}(x) = \frac{5}{6}f_{\mathbf{SN}_1}(x) - f_{\mathbf{SN}_2}(x) = \frac{1}{24} \times k_{29}(x),$$

with

$$k_{29}(x) = 6(\sqrt{x} + 1)\sqrt{2x + 2} - \left[2\sqrt{2x^2 + 2} + 5(\sqrt{x} + 1)^2 \right].$$

In order to prove $k_{29}(x) > 0, \forall x > 0, x \neq 1$, here we shall apply twice the argument given in part 2. Let us consider

$$\begin{aligned} h_{29}(x) &= \left[6(\sqrt{x} + 1)\sqrt{2x + 2} \right]^2 - 2 \left[2\sqrt{2x^2 + 2} + 5(\sqrt{x} + 1)^2 \right]^2 \\ &= 39x^2 + 41x^{3/2} + 3\sqrt{x}(\sqrt{x} - 1)^2 + 41\sqrt{x} + 39 \\ &\quad - 20(\sqrt{x} + 1)^2\sqrt{2x^2 + 2}. \end{aligned}$$

Applying again over $h_{29}(x)$ the argument given in part 2, we get

$$\begin{aligned} h_{29a}(x) &= \left[39x^2 + 41x^{3/2} + 3\sqrt{x}(\sqrt{x} - 1)^2 + 41\sqrt{x} + 39 \right]^2 \\ &\quad - \left[20(\sqrt{x} + 1)^2\sqrt{2x^2 + 2} \right]^2 \\ &= (\sqrt{x} - 1)^4 (721x^2 + 3116x^{3/2} + 4806x + 3116\sqrt{x} + 721)^2. \end{aligned}$$

Since $h_{29a}(x) > 0$ giving $h_{29}(x) > 0, \forall x > 0, x \neq 1$ and consequently, we have $k_{29}(x) > 0, \forall x > 0, x \neq 1$. Thus we have $f_{\mathbf{SN}_1\text{-}\mathbf{SN}_2}(x) > 0, \forall x > 0, x \neq 1$, thereby proving the required result.

30. **For $\mathbf{D}_{\mathbf{SN}_3} \leq \frac{8}{9}\mathbf{D}_{\mathbf{SN}_1}$** : This result is already appearing in (11).
31. **For $\mathbf{D}_{\mathbf{SN}_1} \leq \frac{1}{2}\mathbf{D}_{\mathbf{P}_6\mathbf{G}}$** : By considering the function $g_{\mathbf{SN}_1\text{-}\mathbf{P}_6\mathbf{G}}(x) = f''_{\mathbf{SN}_1}(x)/f''_{\mathbf{P}_6\mathbf{G}}(x)$, we get $\beta_{\mathbf{SN}_1\text{-}\mathbf{P}_6\mathbf{G}} = g_{\mathbf{SN}_1\text{-}\mathbf{P}_6\mathbf{G}}(1) = \frac{1}{2}$, where $f''_{\mathbf{SN}_1}(x)$ and $f''_{\mathbf{P}_6\mathbf{G}}(x)$ are as given by (37) and (20) respectively. We can write

$$\frac{1}{2}D_{\mathbf{P}_6\mathbf{G}} - D_{\mathbf{SN}_1} = b f_{\mathbf{P}_6\mathbf{G}\text{-}\mathbf{SN}_1} \left(\frac{a}{b} \right),$$

where

$$f_{\mathbf{P}_6\mathbf{G}\text{-}\mathbf{SN}_1}(x) = \frac{1}{2}f_{\mathbf{P}_6\mathbf{G}}(x) - f_{\mathbf{SN}_1}(x) = \frac{k_{31}(x)}{4(x + 1)},$$

with

$$k_{31}(x) = 3x^2 + 2x + 3 - 2(x+1)\sqrt{2x^2+2}.$$

Now we shall show that $k_{31}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{31}(x) = (3x^2 + 2x + 3)^2 - \left[2(x+1)\sqrt{2x^2+2}\right]^2$$

After simplifications we have

$$h_{31}(x) = (x-1)^4.$$

Since $h_{31}(x) > 0$, giving $k_{31}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_6G_SN_1}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

32. **For $D_{SN_1} \leq \frac{3}{4}D_{SG}$** : This result is already appearing in (11).

33. **For $D_{SG} \leq \frac{4}{5}D_{P_5H}$** : By considering the function $g_{SG_P_5H}(x) = f''_{SG}(x)/f''_{P_5H}(x)$, we get $\beta_{SG_P_5H} = g_{SG_P_5H}(1) = \frac{4}{5}$, where $f''_{SG}(x)$ and $f''_{P_5H}(x)$ are as given by (42) and (29) respectively. We can write

$$\frac{4}{5}D_{P_5H} - D_{SG} = b f_{P_5H_SG}\left(\frac{a}{b}\right),$$

where

$$f_{P_5H_SG}(x) = \frac{4}{5}f_{P_5H}(x) - f_{SG}(x) = \frac{k_{33}(x)}{4(x+1)},$$

with

$$k_{33}(x) = 2 \left(4x^3 + 5x^{5/2} + 11x^2 + 3x(\sqrt{x}-1)^2 + 11x + 5\sqrt{x} + 4 \right) - 5(x+1)(\sqrt{x}+1)^2\sqrt{2x^2+2}.$$

Now we shall show that $k_{33}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{33}(x) = \left[2 \left(4x^3 + 5x^{5/2} + 11x^2 + 3x(\sqrt{x}-1)^2 + 11x + 5\sqrt{x} + 4 \right) \right]^2 - \left[5(x+1)(\sqrt{x}+1)^2\sqrt{2x^2+2} \right]^2.$$

After simplifications we have

$$h_{33}(x) = 2(\sqrt{x}-1)^4 \left(\begin{array}{l} 7x^4 + 8x^{7/2} + 64x^3 + 120x^{5/2} + \\ + 242x^2 + 120x^{3/2} + 64x + 8\sqrt{x} + 7 \end{array} \right).$$

Since $h_{33}(x) > 0$, giving $k_{33}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_5H_SG}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

Parts 27-33 prove the sequences of inequalities appearing in (44).

34. **For $\mathbf{D}_{P_6P_1} \leq \frac{16}{9}\mathbf{D}_{P_5P_2}$** : By considering the function $g_{P_6P_1-P_5P_2}(x) = f''_{P_6P_1}(x)/f''_{P_5P_2}(x)$, we get $\beta_{P_6P_1-P_5P_2} = g_{P_6P_1-P_5P_2}(1) = \frac{16}{9}$, where $f''_{P_6P_1}(x)$ and $f''_{P_5P_2}(x)$ are as given by (23) and (31) respectively. We can write

$$\frac{16}{9}D_{P_5P_2} - D_{P_6P_1} = b f_{P_5P_2-P_6P_1}\left(\frac{a}{b}\right),$$

where

$$\begin{aligned} f_{P_5P_2-P_6P_1}(x) &= \frac{16}{9}f_{P_5P_2}(x) - f_{P_6P_1}(x) \\ &= \frac{(\sqrt{x}-1)^4 \left(7x^5 + 10x^{9/2} + 23x^4 + 64x^{7/2} + 142x^3 + \right. \\ &\quad \left. + 180x^{5/2} + 142x^2 + 64x^{3/2} + 23x + 10\sqrt{x} + 7 \right)}{9(x^2+1)(x^3+1)(\sqrt{x}+1)^2}. \end{aligned}$$

Since $f_{P_5P_2-P_6P_1}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

35. **For $\mathbf{D}_{P_5P_1} \leq \frac{13}{12}\mathbf{D}_{P_6P_2}$** : By considering the function $g_{P_5P_1-P_6P_2}(x) = f''_{P_5P_1}(x)/f''_{P_6P_2}(x)$, we get $\beta_{P_5P_1-P_6P_2} = g_{P_5P_1-P_6P_2}(1) = \frac{13}{12}$, where $f''_{P_5P_1}(x)$ and $f''_{P_6P_2}(x)$ are as given by (31) and (22) respectively. We can write

$$\frac{13}{12}D_{P_6P_2} - D_{P_5P_1} = b f_{P_6P_2-P_5P_1}\left(\frac{a}{b}\right),$$

where

$$\begin{aligned} f_{P_6P_2-P_5P_1}(x) &= \frac{13}{12}f_{P_6P_2}(x) - f_{P_5P_1}(x) \\ &= \frac{(\sqrt{x}-1)^4 \left(x^5 + 30x^{9/2} + 89x^4 + 152x^{7/2} + 181x^3 + \right. \\ &\quad \left. + 190x^{5/2} + 181x^2 + 152x^{3/2} + 89x + 30\sqrt{x} + 1 \right)}{12(x^2+1)(x^3+1)(\sqrt{x}+1)^2}. \end{aligned}$$

Since $f_{P_6P_2-P_5P_1}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

36. **For $\mathbf{D}_{P_5P_1} \leq \frac{13}{9}\mathbf{D}_{P_5P_2}$** : By considering the function $g_{P_5P_1-P_5P_2}(x) = f''_{P_5P_1}(x)/f''_{P_5P_2}(x)$, we get $\beta_{P_5P_1-P_5P_2} = g_{P_5P_1-P_5P_2}(1) = \frac{13}{9}$, where $f''_{P_5P_1}(x)$ and $f''_{P_5P_2}(x)$ are as given by (32) and (31) respectively. We can write

$$\frac{13}{9}D_{P_5P_2} - D_{P_5P_1} = b f_{P_5P_2-P_5P_1}\left(\frac{a}{b}\right),$$

where

$$\begin{aligned} f_{P_5P_2-P_5P_1}(x) &= \frac{13}{9}f_{P_5P_2}(x) - f_{P_5P_1}(x) \\ &= \frac{(\sqrt{x}-1)^4 \left(4x^5 + 16x^{9/2} + 44x^4 + 88x^{7/2} + 139x^3 + \right. \\ &\quad \left. + 162x^{5/2} + 139x^2 + 88x^{3/2} + 16\sqrt{x} + 44x + 4 \right)}{9(x^2+1)(x^3+1)(\sqrt{x}+1)^2}. \end{aligned}$$

Since $f_{P_5P_2-P_5P_1}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

37. **For $\mathbf{D}_{P_6P_2} \leq \frac{12}{7}\mathbf{D}_{P_5P_3}$** : By considering the function $g_{P_6P_2-P_5P_3}(x) = f''_{P_6P_2}(x)/f''_{P_5P_3}(x)$, we get $\beta_{P_6P_2-P_5P_3} = g_{P_6P_2-P_5P_3}(1) = \frac{12}{7}$, where $f''_{P_6P_2}(x)$ and $f''_{P_5P_3}(x)$ are as given by (22) and (30) respectively. We can write

$$\frac{12}{7}D_{P_5P_3} - D_{P_6P_2} = b f_{P_5P_3-P_6P_2}\left(\frac{a}{b}\right),$$

where

$$\begin{aligned} f_{P_5P_3-P_6P_2}(x) &= \frac{16}{9}f_{P_5P_3}(x) - f_{P_6P_2}(x) \\ &= \frac{(\sqrt{x}-1)^4 \left(5x^4 + x^{7/2} + 17x^3 + 22x^{5/2} + 38x^2 + 22x^{3/2} + 17x + \sqrt{x} + 5 \right)}{7(x^2+1)(x^3+1)(\sqrt{x}+1)^2}. \end{aligned}$$

Since $f_{P_5P_3-P_6P_2}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

38. **For $\mathbf{D}_{P_5P_2} \leq \frac{9}{7}\mathbf{D}_{P_5P_3}$** : By considering the function $g_{P_5P_2-P_5P_3}(x) = f''_{P_5P_2}(x)/f''_{P_5P_3}(x)$, we get $\beta_{P_5P_2-P_5P_3} = g_{P_5P_2-P_5P_3}(1) = \frac{9}{7}$, where $f''_{P_5P_2}(x)$ and $f''_{P_5P_3}(x)$ are as given by (31) and (30) respectively. We can write

$$\frac{9}{7}D_{P_5P_3} - D_{P_5P_2} = b f_{P_5P_3-P_5P_2}\left(\frac{a}{b}\right),$$

where

$$\begin{aligned} f_{P_5P_3-P_5P_2}(x) &= \frac{9}{7}f_{P_5P_3}(x) - f_{P_5P_2}(x) \\ &= \frac{(\sqrt{x}-1)^4 \left(2x^3 + 6x^{5/2} + 16x^2 + 21x^{3/2} + 16x + 6\sqrt{x} + 2 \right)}{7(x^2+1)(x-\sqrt{x}+1)(\sqrt{x}+1)^2}. \end{aligned}$$

Since $f_{P_5P_3-P_5P_2}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

39. **For $\mathbf{D}_{P_5P_3} \leq \frac{14}{9}\mathbf{D}_{P_6N_2}$** : By considering the function $g_{P_5P_3-P_6N_2}(x) = f''_{P_5P_3}(x)/f''_{P_6N_2}(x)$, we get $\beta_{P_5P_3-P_6N_2} = g_{P_5P_3-P_6N_2}(1) = \frac{14}{9}$, where $f''_{P_5P_3}(x)$ and $f''_{P_6N_2}(x)$ are as given by (30) and (17) respectively. We can write

$$\frac{14}{9}D_{P_6N_2} - D_{P_5P_3} = b f_{P_6N_2-P_5P_3}\left(\frac{a}{b}\right),$$

where

$$\begin{aligned} f_{P_6N_2-P_5P_3}(x) &= \frac{14}{9}f_{P_6N_2}(x) - f_{P_5P_3}(x) \\ &= \frac{(\sqrt{x}-1)^4 \left(x^6 + 268x^{11/2} + 1143x^5 + 782x^{9/2} + 2597x^4 + 2014x^{7/2} + 4478x^3 + 2014x^{5/2} + 2597x^2 + 782x^{3/2} + 1143x + 268\sqrt{x} + 1 \right)}{18(x^3+1)(\sqrt{x}+1)^2}. \end{aligned}$$

Since $f_{P_6N_2-P_5P_3}(x) > 0, \forall x > 0, x \neq 1$, hence proving the required result.

40. **For $\mathbf{D}_{P_6N_2} \leq \frac{9}{4}\mathbf{D}_{P_6S}$** : By considering the function $g_{P_6N_2-P_6S}(x) = f''_{P_6N_2}(x)/f''_{P_6S}(x)$, we get $\beta_{P_6N_2-P_6S} = g_{P_6N_2-P_6S}(1) = \frac{9}{4}$, where $f''_{P_6N_2}(x)$ and $f''_{P_6S}(x)$ are as given by (17) and (16) respectively. We can write

$$\frac{9}{4}D_{P_6S} - D_{P_6N_2} = b f_{P_6S-P_6N_2}\left(\frac{a}{b}\right),$$

where

$$f_{P_6S-P_6N_2}(x) = \frac{9}{4}f_{P_6S}(x) - f_{P_6N_2}(x) = \frac{k_{40}(x)}{8(x+1)},$$

with

$$k_{40}(x) = 2\sqrt{2x+2}(\sqrt{x}+1)(x+1) + 10x^2 + 10 - 9(x+1)\sqrt{2x^2+2}.$$

In order to prove $k_{40}(x) > 0$, $\forall x > 0$, $x \neq 1$, we shall apply twice the argument given in part 2. Let us consider

$$\begin{aligned} h_{40}(x) &= \left[2\sqrt{2x+2}(\sqrt{x}+1)(x+1) + 10x^2 + 10\right]^2 - \left[9(x+1)\sqrt{2x^2+2}\right]^2 \\ &= 40\sqrt{2x+2}(\sqrt{x}+1)(x^2+1)(x+1) \\ &\quad - \left[46x^4 + 8(x+1)^3(\sqrt{x}-1)^2 + 260x^3 + 28x^2 + 260x + 46\right]. \end{aligned}$$

Applying again over $h_{40}(x)$ the argument given in part 2, we get

$$\begin{aligned} h_{40a}(x) &= \left[40\sqrt{2x+2}(\sqrt{x}+1)(x^2+1)(x+1)\right]^2 \\ &\quad - \left[46x^4 + 8(x+1)^3(\sqrt{x}-1)^2 + 260x^3 + 28x^2 + 260x + 46\right]^2 \\ &= 4(\sqrt{x}-1)^4 \left(\begin{aligned} &71 + 2316\sqrt{x} + 11960x^{5/2} + 4090x + 11960x^{7/2} + \\ &+ 11180x^{3/2} + 2316x^{11/2} + 11180x^{9/2} + 12021x^4 + \\ &+ 12021x^2 + 6004x^3 + 4090x^5 + 71x^6 \end{aligned} \right). \end{aligned}$$

Since $h_{40a}(x) > 0$ giving $h_{40}(x) > 0$, $\forall x > 0$, $x \neq 1$ and consequently, we have $k_{40}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{P_6S-P_6N_2}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

41. **For $\mathbf{D}_{P_6S} \leq \mathbf{D}_{AG}$** : By considering the function $g_{P_6S-AG}(x) = f''_{P_6S}(x)/f''_{AG}(x)$, we get $\beta_{P_6S-AG} = g_{P_6S-AG}(1) = 1$, where $f''_{P_6S}(x)$ and $f''_{AG}(x)$ are as given by (16) and (17) respectively. We can write

$$D_{AG} - D_{P_6S} = b f_{AG-P_6S}\left(\frac{a}{b}\right),$$

where

$$f_{AG-P_6S}(x) = f_{AG}(x) - f_{P_6S}(x) = \frac{k_{41}(x)}{2(x+1)},$$

with

$$k_{41}(x) = \sqrt{2x^2 + 2}(x + 1) - [(x - 1)^2 + 2(x + 1)\sqrt{x}].$$

Now we shall show that $k_{41}(x) > 0$, $\forall x > 0$, $x \neq 1$. Let us consider

$$h_{41}(x) = \left[\sqrt{2x^2 + 2}(x + 1) \right]^2 - [(x - 1)^2 + 2(x + 1)\sqrt{x}]^2$$

After simplifications we have

$$h_{41}(x) = (\sqrt{x} + 1)^2 (\sqrt{x} - 1)^6.$$

Since $h_{41}(x) > 0$, giving $k_{41}(x) > 0$, $\forall x > 0$, $x \neq 1$. Thus we have $f_{AG-P_6S}(x) > 0$, $\forall x > 0$, $x \neq 1$, thereby proving the required result.

Parts 34-41 prove the sequences of inequalities appearing in (45).

□

Remark 3.2. (i) In view of Theorem 3.1, we have the following improvement over the inequalities (11):

$$D_{SA} \leq \left\{ \begin{array}{l} \frac{1}{3}D_{SH} \leq \frac{1}{2}D_{AH} \leq 4D_{N_2N_1} \leq \frac{4}{3}D_{N_2G} \leq D_{AG} \leq 4D_{AN_2} \\ \left\{ \begin{array}{l} \frac{4}{5}D_{SN_2} \\ \frac{3}{4}D_{SN_3} \end{array} \right\} \leq \frac{2}{3}D_{SN_1} \leq \frac{1}{2}D_{SG} \end{array} \right\}.$$

(ii) The results appearing Parts 1-41 bring us some very interesting measures given by

$$V_k(a, b) = b f_k(a/b), \quad k = 1, 2, 3, 4,$$

where

$$f_1(x) = \frac{(x + 1)^2 (x - 1)^4}{(x^3 + 1)(x^2 + 1)},$$

$$f_2(x) = \frac{(\sqrt{x} - 1)^2}{x + 1},$$

$$f_3(x) = \frac{\sqrt{x}(\sqrt{x} - 1)^4}{(x + 1)(\sqrt{x} + 1)^2}$$

and

$$f_4(x) = \frac{(\sqrt{x} - 1)^4 [(\sqrt{x} - 1)^2 + \sqrt{x}]}{12(x + 1)}.$$

The measures $V_1(a, b)$ is due to part 1, the measure $V_2(a, b)$ is due to parts 6, 8, 13, 23 and 24, the measure $V_3(a, b)$ is due to parts 9 and 16 and the measure $V_4(a, b)$ is due to parts 14 and 15.

4 Connections with Information Measures

$$\Gamma_n = \left\{ P = (p_1, p_2, \dots, p_n) \left| p_i > 0, \sum_{i=1}^n p_i = 1 \right. \right\}, \quad n \geq 2,$$

be the set of all complete finite discrete probability distributions. For all $P, Q \in \Gamma_n$, the following inequalities are already proved by author [9]:

$$\frac{1}{2}D_{AH} \leq I \leq 4D_{N_2N_1} \leq \frac{4}{3}D_{N_2G} \leq D_{AG} \leq 4D_{AN_2} \leq \frac{1}{8}J \leq T \leq \frac{1}{16}\Psi, \quad (46)$$

where

$$I(P||Q) = \frac{1}{2} \left[\sum_{i=1}^n p_i \ln \left(\frac{2p_i}{p_i + q_i} \right) + \sum_{i=1}^n q_i \ln \left(\frac{2q_i}{p_i + q_i} \right) \right], \quad (47)$$

$$J(P||Q) = \sum_{i=1}^n (p_i - q_i) \ln \left(\frac{p_i}{q_i} \right), \quad (48)$$

$$T(P||Q) = \sum_{i=1}^n \left(\frac{p_i + q_i}{2} \right) \ln \left(\frac{p_i + q_i}{2\sqrt{p_i q_i}} \right) \quad (49)$$

and

$$\Psi(P||Q) = \sum_{i=1}^n \frac{(p_i - q_i)^2 (p_i + q_i)}{p_i q_i}. \quad (50)$$

The measures $I(P||Q)$, $J(P||Q)$ and $T(P||Q)$ are the respectively, the well-know *Jensen-Shannon divergence*, *J-divergence* and *arithmetic and geometric mean divergence*. The measure $\Psi(P||Q)$ is *symmetric chi-square divergence*. For detailed study on these measures refer to Taneja [8, 9, 11]. Moreover, $D_{AH}(P||Q) = \frac{1}{2}\Delta(P||Q)$ and $D_{AG}(P||Q) = h(P||Q)$, where $\Delta(P||Q)$ and $h(P||Q)$ are the well-known *triangular's* and *Hellinger's discriminations* respectively.

In the theorem below we shall make connections of the classical divergence measures given in (47)-(50) with the inequalities given in (43). Moreover the theorem below unifies the inequalities (43) and (46).

Theorem 4.1. *The following inequalities hold:*

$$\left\{ \frac{\frac{2}{3}D_{P_5H}}{\frac{2}{3}D_{AP_4}} \leq I \right\} \leq 4D_{N_2N_1} \leq \frac{4}{3}D_{N_2G} \leq D_{AG} \leq 4D_{AN_2} \leq \frac{2}{3}D_{P_5G} \leq \left\{ \frac{D_{P_5N_1}}{\frac{1}{8}J} \leq \frac{6}{5}D_{P_5N_3} \leq \frac{4}{3}D_{P_5N_2} \leq 2D_{P_5A} \right\} \leq T \leq \frac{1}{16}\Psi.$$

Proof. In view of the inequalities given in (43) and (46), it sufficient to show the following three inequalities:

- (i) $D_{AP_4} \leq \frac{3}{2}I$;
- (ii) $D_{P_5G} \leq \frac{3}{16}J$;
- (iii) $D_{P_5A} \leq \frac{1}{2}T$.

Since in each part of the above expressions we have logarithmic form, we shall apply a different approach to show the above three inequalities.

(i) For $D_{AP_4} \leq \frac{3}{2}I$: Let us consider

$$g_{AP_4-I}(x) = \frac{f''_{AP_4}(x)}{f''_I(x)} = \frac{12x(x+1)}{\sqrt{x}(\sqrt{x}+1)^4}, \quad x \neq 1, \quad x > 0.$$

Calculating the first order derivative of the function $g_{AP_4-I}(x)$ with respect to x , one gets

$$g'_{AP_4-I}(x) = -\frac{6(x^{3/2} - 3x + 3\sqrt{x} - 1)}{\sqrt{x}(\sqrt{x}+1)^5} = -\frac{6(\sqrt{x}-1)^3}{\sqrt{x}(\sqrt{x}+1)^5} \begin{cases} > 0, & x < 1 \\ < 0, & x > 1 \end{cases} \quad (51)$$

In view of (51), we conclude that the function $g_{AP_4-I}(x)$ is increasing in $x \in (0, 1)$ and decreasing in $x \in (1, \infty)$, and hence

$$\beta_{AP_4-I} = \sup_{x \in (0, \infty)} g_{AP_4-I}(x) = g_{AP_4-I}(1) = \frac{3}{2}. \quad (52)$$

By the application of Lemma 3.1 with (52) we get the required result.

(ii) For $D_{P_5G} \leq \frac{3}{16}J$: Let us consider

$$g_{P_5G-J}(x) = \frac{f''_{P_5G}(x)}{f''_J(x)} = \frac{3\sqrt{x}(4x^{3/2} + 4\sqrt{x} + x^2 - 2x + 1)}{4(x+1)(\sqrt{x}+1)^4}.$$

Calculating the first order derivative of the function $g_{P_5G-J}(x)$ with respect to x , one gets

$$g'_{P_5G-J}(x) = -\frac{3(\sqrt{x}-1)^3(x^2 + 8x^{3/2} + 6x + 8\sqrt{x} + 1)}{8\sqrt{x}(\sqrt{x}+1)^5(x+1)^2} \begin{cases} > 0, & x < 1 \\ < 0, & x > 1 \end{cases} \quad (53)$$

In view of (53), we conclude that the function $g_{P_5G-J}(x)$ is increasing in $x \in (0, 1)$ and decreasing in $x \in (1, \infty)$, and hence

$$\beta_{P_5G-J} = \sup_{x \in (0, \infty)} g_{P_5G-J}(x) = g_{P_5G-J}(1) = \frac{3}{2}. \quad (54)$$

By the application of Lemma 3.1 with (54) we get the required result.

(iii) For $\mathbf{D}_{P_5A} \leq \frac{1}{2}\mathbf{T}$: Let us consider

$$g_{P_5A\text{-}T}(x) = \frac{f''_{P_5A}(x)}{f''_T(x)} = \frac{2\sqrt{x}(4x^{3/2} + 4\sqrt{x} + x^2 - 6x + 1)(x+1)}{(x^2+1)(\sqrt{x}+1)^4}.$$

Calculating the first order derivative of the function $g_{P_5A\text{-}T}(x)$ with respect to x , one gets

$$g'_{P_5A\text{-}T}(x) = -\frac{(\sqrt{x}-1)^3 \left(\frac{8\sqrt{x}(x^2+1)(\sqrt{x}-1)^2}{+x^4+14x^3+10x^2+14x+1} \right)}{\sqrt{x}(\sqrt{x}+1)^5(x^2+1)^2} \begin{cases} > 0, & x < 1 \\ < 0, & x > 1 \end{cases} \quad (55)$$

In view of (55), we conclude that the function $g_{P_5A\text{-}T}(x)$ is increasing in $x \in (0, 1)$ and decreasing in $x \in (1, \infty)$, and hence

$$\beta_{P_5A\text{-}T} = \sup_{x \in (0, \infty)} g_{P_5A\text{-}T}(x) = g_{P_5A\text{-}T}(1) = \frac{1}{2}. \quad (56)$$

By the application of Lemma 3.1 with (56) we get the required result.

Parts (i)-(iii) completes the proof of the Theorem 4.1. $\square$

Here we have referred Lemma 3.1, but its extension for the probability distributions is already proved in [9, 10].

References

- [1] P. CZINDER and Z. PALES, Local monotonicity properties of two-variable Gini means and the comparison theorem revisited, *J. Math. Anal. Appl.*, **301**(2005), 427-438.
- [2] CHAO-PING CHEN, Asymptotic Representations for Stolarsky, Gini and the Generalized Muirhead Means, *RGMIA COLLECTION*, **11**(4)(2008), 1-13.
- [3] C. GINI, Di una formula compressiva delle medie, *Metron*, **13**(1938) 322.
- [4] D. H. LEHMER, On the compounding of certain means, *J. Math. Anal. Appl.*, **36**(1971), 183-200.
- [5] J. SÁNDER, A Note on Gini Mean, *General Mathematics*, **12**(4)(2004), 17-21.
- [6] H. N. SHI, J. ZHANG and DA-MAO LI, Schur-Geometric Convexity for Difference of Means, *Applied Mathematics E-Notes*, **10**(2010), 275-284

- [7] S. SIMIĆ, A Simple Proof of Monotonicity for Stolarsky and Gini Means, *Kragujevac J. Math.*, **32**(2009), 75-79.
 - [8] I.J. TANEJA, New Developments in Generalized Information Measures, Chapter in: *Advances in Imaging and Electron Physics*, Ed. P.W. Hawkes, **91**(1995), 37-136.
 - [9] I.J. TANEJA, On Symmetric and Nonsymmetric Divergence Measures and Their Generalizations, Chapter in: *Advances in Imaging and Electron Physics*, **138**(2005), 177-250.
 - [10] I.J. TANEJA, Refinement Inequalities Among Symmetric Divergence Measures, *The Australian Journal of Mathematical Analysis and Applications*, **2**(1)(2005), Art. 8, pp. 1-23.
 - [11] I.J. TANEJA, Refinement of Inequalities among Means, *Journal of Combinatorics, Information and Systems Sciences*, **31**(2006), 357-378.
-